

Anti-Perlecan (bovine, human)**Mouse monoclonal antibody**

Subclass: IgG1/k

CAT. NO.

CSI 001-74

Clone:A74

SPECIFICITY CSI 001-74 is highly specific for perlecan. There is no evidence for cross-reactivity with other connective tissue proteins (vitronectin, fibronectin, elastin, collagen, laminin).

IMMUNOGEN Bovine corneal endothelial cells

TESTED APPLICATIONS ELISA, WB, IHC-F, IHC-P, IP, AP

SPECIES REACTIVITY (POSITIVE) Bovine, human

SPECIES REACTIVITY (NEGATIVE) Not determined

EPITOPE SPECIFICITY Epitope is located in domain V

PRESENTATION

Content: Available in 200 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.

Preparation: Protein-A purified

Form: Liquid

Solvent: 0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide

Storage: 4-8°C without exposure to light. No precautions necessary during handling.

APPLICATION

ELISA: CSI 001-74 can be used in ELISA

WB: In Western blotting a dilution guideline of 1/100 has proved successful (1).

IHC: CSI 001-74 can be used in immunostaining of frozen PLP-fixed sections of bovine and human tissues.

AP: CSI 001-74 can be used in affinity chromatography, in conjunction with CSI 001-71 to separate recombinant domain I from full length perlecan (1).

IP: CSI 001-74 can be used in immunoprecipitation

TARGET

Perlecan is an extracellular matrix proteoglycan. It has a large core protein of 400-450 kDa and is often produced with heparan sulfate side chains. Perlecan is found in basement membranes where it contributes to the permeability characteristics, serves as a substrate for vascular cells and binds growth factors involved in vascular remodelling.

REFERENCES

1. Whitelock JM, Murdoch AD, Iozzo RV, Underwood PA (1996) The degradation of human endothelial cell-derived perlecan, and release of bound bFGF by stromelysin, plasmin and heparanases. J Biol Chem 271:10079-10086.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.