

**Anti-neutrophil gelatinase-associated lipocalin (human, NGAL)
Mouse monoclonal antibody**

Subclass: IgG1/k

PRODUCT NO.

ABS 038-26

Clone: 26

PRESENTATION

Preparation: Protein-A/G purified

Content: Available in 200 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.

Solvent: 0.01 M phosphate buffer, pH 7.4, with 0.5 M NaCl and 15 mM sodium azide

Storage: 4-8°C without exposure to light. No precautions necessary during handling.

ANTIGEN

Neutrophil gelatinase-associated lipocalin (NGAL; also called lipocalin 2, siderocalin and neutrophil lipocalin) is a member of the lipocalin family of proteins which bind and transport small lipophilic molecules. NGAL is released by activated neutrophils and occurs as 25-kDa glycosylated single protein chain monomers, which form dimers and small amounts of higher oligomers, as well as complexes with matrix metalloproteinase 9 (MMP-9; gelatinase B) (1). Low level expression of NGAL in a variety of epithelia may be increased in inflammation or cancers (2). Expression of NGAL in the kidney is dramatically increased by acute kidney injury (3).

IMMUNOGEN

Recombinant human NGAL adsorbed onto aluminum hydroxide gel

SPECIFICITY

ABS 038-26 binds human NGAL and does not cross-react with NGAL from other mammalian species tested

EPI TOPE SPECIFICITY

Differs from those of HYB 211-01, ABS 038-14, ABS 038-15 and ABS 038-23

REACTIVITY

ABS 038-26 binds human NGAL in solution and coated onto a solid phase.

ABS 038-26 (as capture antibody) forms sandwich pairs with HYB 211-01, ABS 038-14, ABS 038-15 and ABS 038-23 (as biotinylated detection antibodies) for measuring human NGAL. In Western blotting a dilution guideline of 1/1000 has proved successful.

CULTURE MEDIUM

RPMI 1640 with 2-10% fetal calf serum

FUSION PARTNER

SP2mIL6

IMMUNIZATION

Female NMRIxBALB/c mice immunized by intraperitoneal injection

APPLICATION

Method	Usability	References
ELISA	Yes	
Immunoblotting	Yes	
Immunohistochemistry	Not determined	

REFERENCES

1. Kjeldsen L, Johnsen AH, Sengeløv H, Borregaard N (1993) Isolation and primary structure of NGAL, a novel protein associated with human neutrophil gelatinase. *J Biol Chem* 268:10425-10432.
2. Nielsen BS, Borregaard N, Bundgaard JR, Timshel S, Sehested M, Kjeldsen L (1996) Induction of NGAL synthesis in epithelial cells of human colorectal neoplasia and inflammatory bowel diseases. *Gut* 38:414-420.
3. Matthaeus T, Schulze-Lohoff E, Ichimura T, Weber M, Andreucci M, Park KM, Alessandrini A, Bonventre JV (2001) Co-regulation of neutrophil gelatinase-associated lipocalin and matrixmetalloproteinase-9 in the postischemic rat kidney. *J Am Soc Nephrol* 12:787A.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.