

**Anti-Complement component C3d (human)
 Mouse monoclonal antibody**

Subclass: IgG₁/k

PRODUCT NO.

HYB 030-08

Clone: 6F6

PRESENTATION

Preparation: Protein-A/G purified

Content: Available in 200 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.

Solvent: 0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide

Storage: 4-8°C without exposure to light. No precautions necessary during handling.

ANTIGEN

C3 is a key component of the complement system since classical and alternative activation pathways merge at the C3 activation step when C3 is split into C3a and C3b. The molecular mass of C3 is 185 kDa and it consists of two chains (110 kDa and 75 kDa) held together by disulfide bonds (1,2).

IMMUNOGEN

C3 isolated from human plasma adsorbed onto aluminum hydroxide gel

SPECIFICITY

HYB 030-08 is specific for human C3d, a fragment of the alpha chain (3)

EPI TOPE SPECIFICITY

Epitope specificity differs from that of HYB 005-01, HAV 003-05 and HAV 004-01 but slightly overlap as determined by inhibition ELISA.

REACTIVITY

HYB 030-08 reacts strongly with C3. A strong reaction is seen in ELISA with C3 directly coated onto the microtiterwell, or when used as a biotinylated detection antibody in sandwich ELISA in combination with a polyclonal antibody against C3 (e.g. DAKO A 0062).

In Western blotting after SDS-PAGE, HYB 030-08 reacts with C3 in reduced as well as nonreduced forms.

CULTURE MEDIUM

Dulbecco's modified Eagle's medium with 10% fetal calf serum

FUSION PARTNER

X63-Ag8.653

IMMUNIZATION

Female CF1 x BALB/c mice immunized by intraperitoneal injection

APPLICATION

Method	Usability	References
ELISA	Yes	
Immunoblotting	Yes	
Immunohistochemistry	Not determined	

REFERENCES

1. Law SKA, Reid KBM (1988) Complement. In: In Focus (Ed. Male D) IRL Press: Oxford.
2. Morley BJ and Walport MJ (2000) The Complement FactsBook. Academic Press, London, UK.
3. Nagar B, Jones RG, Diefenbach RJ, Isenman DE, Rini JM (1998) X-ray crystal structure of C3d: a C3 fragment and ligand for complement receptor 2. Science 280:1277-1281.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.