

**Anti-Mannan-binding lectin (human, MBL)
 Mouse monoclonal antibody**

Subclass: IgG1/k

PRODUCT NO.

HYB 131-11

Clone: 11C9

PRESENTATION

Preparation: Protein-A/G purified

Content: Available in 200 µL and 1 mL size. 1 mg/mL +/- 15%. See Certificate of Analysis for details.

Solvent: 0.01 M phosphate buffer, pH 7.4, containing 0.5 M NaCl and 15 mM sodium azide

Storage: 4-8°C without exposure to light. No precautions necessary during handling.

ANTIGEN

Mannan-binding lectin (MBL), also called mannanose-binding lectin or protein, belongs to the C-type family of collectins, showing calcium-dependent binding to certain sugars (1). It consists of oligomers of triple-chain subunits and its binding and complement activating activities depend on its normal oligomerization. On binding to mannan-like microbial surface carbohydrates, MBL activates the complement system by means of its own lectin pathway, depending on the MBL-associated serine proteases (MASPs). Because of the presence of different structural and promoter alleles in the population, 12% or more of the population have low concentrations (<50ng/mL) of normally oligomerized, functional MBL in plasma or serum (2).

IMMUNOGEN

MBL purified from human donor plasma and adsorbed onto aluminum hydroxide

SPECIFICITY

HYB 131-11 is specific for MBL from human serum or plasma.

EPI TOPE SPECIFICITY

The epitope is thought to be on the carbohydrate recognition domain and differs from that of HYB 131-01 and HYB 131-10.

REACTIVITY

HYB 131-11 reacts strongly with MBL. Strong reaction is seen in ELISA with MBL coated directly onto the microtiter well. In Western blotting HYB 131-11 reacts with human MBL in both its oligomerized state and as single protein chain of 26 kDa. HYB 131-11 can be used to measure MBL in plasma or serum and is specific for oligomerized MBL when used in combination with itself, but not with other MBL antibodies, when it also reacts with poorly oligomerized forms (3).

CULTURE MEDIUM

RPMI 1640 with 10% fetal calf serum

FUSION PARTNER

X63-Ag8.653

IMMUNIZATION

Female CF1 x BALB/c mice immunized by intraperitoneal injection

APPLICATION

Method	Usability	References
ELISA	Yes	
Immunoblotting	Yes	
Immunohistochemistry	Not determined	

REFERENCES

1. Kawasaki N, Kawasaki T, Yamashina I (1983) Isolation and characterization of a mannan-binding protein from human serum. J Biochem (Tokyo) 94:937-947.
2. Steffensen R, Thiel S, Varming K, Jersild C, Jensenius JC (2000) Detection of structural gene mutations and promoter polymorphisms in the mannan-binding lectin (MBL) gene by polymerase chain reaction with sequence-specific primers. J Immunol Methods 241:33-42.
3. Garred P, Larsen F, Madsen HO, Koch C (2003) Mannose-binding lectin deficiency - revisited. Mol Immunol 40:73-84.

CONDITIONS

Unless otherwise marked, all products are for research use only. Not for use in diagnostic procedures. Not for use in human therapeutic applications. For in vitro use or further manufacture only. The information and product are offered without guarantee as the ultimate conditions of use are beyond our control. The foregoing is in lieu of all warranties, expressed or implied, including implied warranties of merchantability and fitness for a particular purpose. In no event shall BioPorto Diagnostics A/S be responsible for loss of profits or indirect consequential losses resulting from use of its products.